

ARMOR

67-F85

RM: 6.7

CHARACTERISTICS


Maturity
Plant Type
Plant Height
Lodging
Shatter
Flower Color
Hilum Color
Pubescence
Pod Color

DISEASE RATINGS

Phytophthora
Phytophthora Gene
Sudden Death
Frogeye Leaf Spot
Stem Canker
Cyst Nematode
Southern RKN

Determinate Medium Resistant NA Purple Black Lt. Tawny Tan Resistant NG Resistant Resistant Resistant Resistant Resistant Resistant Resistant	6.7
Resistant NA Purple Black Lt. Tawny Tan Resistant NG Resistant Resistant Resistant Resistant	Determinate
NA Purple Black Lt. Tawny Tan Resistant NG Resistant Resistant Resistant Resistant	Medium
Purple Black Lt. Tawny Tan Resistant NG Resistant Resistant Resistant Resistant	Resistant
Black Lt. Tawny Tan Resistant NG Resistant Resistant Resistant Resistant	NA
Lt. Tawny Tan Resistant NG Resistant Resistant Resistant Resistant	Purple
Resistant NG Resistant Resistant Resistant	Black
Resistant NG Resistant Resistant Resistant	Lt. Tawny
NG Resistant Resistant Resistant	Tan
NG Resistant Resistant Resistant	
Resistant Resistant Resistant	Resistant
Resistant Resistant	NG
Resistant	Resistant
	Resistant
MR3, MR14	Resistant
	MR3, MR14
Resistant	Resistant

Trait Packages –XTENDFLEX®



- Plant Height -

MEDIUM

— Emergence — **EXCELLENT**

— Chloride Tolerance —
EXCLUDER

Soybean Rating System: There are many ways to evaluate diseases, agronomic characteristics, etc. Since many of these ratings are subjective, an explanation is listed here: Resistant implies no or very slight disease presence when conditions are favorable for development. Moderately Resistant (M. Resistant) implies slight disease presence when conditions are favorable for development. Moderate implies some disease presence when conditions are favorable for development. Moderately Susceptible (M. Susceptible) implies significant disease presence when conditions are favorable for development. Susceptible implies very significant disease presence when conditions are favorable for development.

These are general recommendations based on data taken from breeder, university, company trials and producer field observations and do not constitute a warranty of fitness or guarantee of performance for a particular use. • Double cropping Maturity Group IVs can be extremely risky without adequate irrigation and proper variety selection. • Upright implies a narrow canopy with very little branching. Intermediate implies a wider canopy with some branching. Bushy implies a wide canopy. Imp. Black = Imperfect Black. NG = No Gene. — = Not Available.

Unauthorized propagation, seed multiplication, and/or sale, offering for sale, exchange, delivery, transfer, import, or export of seed of this variety is prohibited. Violators of the U.S. Plant Variety Protection ACT (PVPA) will be subject to proceedings under federal and state law and may be liable for money damages. Seed of varieties protected or applied for under 1994 amendments to the PVPA cannot be sold for reproductive purposes without permission from the owner of this variety.

Ratings are general. Soil and environment may adversely affect performance



PROUD MEMBER OF U.S. AGRICULTURE

